

Non-destructive discrimination of multiparticle cluster states for quantum computation

Srерaman Muralidharan
*Loyola College, Chennai - 600034, India**

Sakshi Jain
Indian Institute of Technology-Bombay, Mumbai - 400080, India†

Sriram Prasath E.‡
Maulana Azad National Institute of Technology, Bhopal - 462051, India

Prasanta K. Panigrahi§
*Indian Institute of Science Education and Research Kolkata, Mohanpur Campus,
BCKV Campus Main Office, Mohanpur - 741252, India and
Physical Research Laboratory, Navrangpura, Ahmedabad - 380 009, India*

Discriminating between orthogonal quantum systems without destroying their entanglement is of interest to quantum computation and communication. In this paper, we explicate the schemes for the non-destructive discrimination (NDD) of 16 orthogonal four qubit cluster states and 32 orthogonal five qubit cluster states respectively. This technique will find its applications in various branches of quantum information theory.

PACS numbers: 03.67.Hk, 03.65.Ud

Keywords: Cluster state, Entanglement, Non-destructive Discrimination

I. INTRODUCTION

Many quantum communication protocols like teleportation [1], state sharing [2, 3] and dense coding [4] require measurement to be performed on entangled bases. For instance, in the dense coding protocol, involving Bell states as a shared entangled resource, two classical bits are encoded into a qubit by the sender and sent to the receiver. The receiver then performs a measurement on the Bell basis and recovers the encoded information. In this process, the entanglement is not disturbed and the same Bell state can be reused for the dense coding of a number of classical bits. Many teleportation protocols also involve Bell measurements. Recently several quantum protocols have been carried out using a special type of graph states known as “cluster states” [5, 6] given by

$$|C_N\rangle = \frac{1}{2^{N/2}} \otimes_{a=1}^N (|0\rangle_a \sigma_z^{a+1} + |1\rangle_a), \quad (1)$$

with $\sigma_z^{N+1} = 1$. These states have also been experimentally realized in various systems [7, 8, 9]. It is well known that they show a strong violation of local reality and are shown to be robust against decoherence [10, 11]. Entanglement exhibited by these states have well been characterized owing to their promising usefulness in Quantum communication protocols [12, 13], one way quantum computation [11] and also for quantum error correction [14]. Two of the present authors [15] have shown them to be useful for perfect teleportation and state sharing of an arbitrary two qubit state. Its superdense coding capacity reaches the Holevo bound. The main hurdle towards the experimental realization of the above mentioned protocols is that most of these schemes involve arbitrary multiqubit measurements [15, 16, 17, 18, 19] which are extremely difficult to realize in laboratory conditions.

In an experimental scenario, since even performing a simple Bell basis measurement is a non trivial task, measurement on a product basis is preferred. Recently, this has been made possible by performing suitable quantum gates which are experimentally realizable. Indeed, this technique has been used for the experimental implementation of

*Electronic address: srерamanm@gmail.com

†Electronic address: sakshi.r.jain@gmail.com

‡Electronic address: sridif@gmail.com

§Electronic address: prasanta@prl.res.in


teleportation in *NMR* [20] and in quantum dots [21] for the teleportation of an arbitrary single qubit state. However, the disadvantage arising out of this technique is that the same resource state cannot be reused for other quantum protocols since the entanglement is destroyed.

Here, we propose an appropriate circuit using which we perform a non-destructive measurement of an entangled basis. Recently, it was shown using an appropriate quantum circuit that four Bell pairs can be discriminated deterministically [22, 23]. The advantage of our scheme over the previous schemes discussed above is that we perform a product basis measurement and yet do not disturb the entanglement of the system. We use this method to discriminate between orthogonal four and five qubit cluster states.


The paper is organized as follows: We initially discuss the circuits for the physical realization of four and five qubit cluster states. The next section deals with the explicit circuits for their NDD. In the last section, we illustrate the usefulness of our circuit for various quantum computational protocols.

II. PHYSICAL REALIZATION

Explicit circuits for the physical realization of the multi-particle cluster states have been studied by two of the present authors [24]. For the sake of completeness, we describe these circuits before we proceed with the non-destructive discrimination of these states. The 16 orthogonal four qubit cluster states can be generated by implementing the following general circuit diagram (Fig. 1) by changing the inputs $|0000\rangle_{1234}$ to a different computational bases.



Similarly, the five-qubit cluster states can be generated by the following circuit (Fig. 2) by changing the inputs



$|00000\rangle_{12345}$ to a different computational bases. Now, we shall proceed with the non-destructive discrimination of these states.

III. NON-DESTRUCTIVE DISCRIMINATION

A. four-qubit Cluster States

The sixteen orthogonal cluster states can be distinguished without disturbing the system by carrying out measurements on four ancilla bits. The four qubit cluster states are made to interact with these ancillas as shown in Fig 3. The outcome of the measurement on the ancillas reveals the respective cluster state as shown in the Table-I:


FIG. 3: Circuit diagram for NDD of $|C_4\rangle$ 

TABLE I: Four-qubit Cluster State discrimination from the ancilla measurements

Ancilla Measurement	Corresponding four-qubit Cluster State
$ 0000\rangle$	$ 0000\rangle + 0011\rangle + 1100\rangle - 1111\rangle$
$ 0001\rangle$	$ 0000\rangle - 0011\rangle + 1100\rangle + 1111\rangle$
$ 0010\rangle$	$ 0001\rangle + 0010\rangle - 1101\rangle + 1110\rangle$
$ 0011\rangle$	$ 0001\rangle - 0010\rangle - 1101\rangle - 1110\rangle$
$ 0100\rangle$	$ 0000\rangle + 0011\rangle - 1100\rangle + 1111\rangle$
$ 0101\rangle$	$ 0000\rangle - 0011\rangle - 1100\rangle - 1111\rangle$
$ 0110\rangle$	$ 0001\rangle + 0010\rangle + 1101\rangle - 1110\rangle$
$ 0111\rangle$	$ 0001\rangle - 0010\rangle + 1101\rangle + 1110\rangle$
$ 1000\rangle$	$ 0100\rangle + 0111\rangle + 1000\rangle - 1011\rangle$
$ 1001\rangle$	$ 0100\rangle - 0111\rangle + 1000\rangle + 1011\rangle$
$ 1010\rangle$	$ 0101\rangle + 0110\rangle - 1001\rangle + 1010\rangle$
$ 1011\rangle$	$ 0101\rangle - 0110\rangle - 1001\rangle - 1010\rangle$
$ 1100\rangle$	$ 0100\rangle + 0111\rangle - 1000\rangle + 1011\rangle$
$ 1101\rangle$	$ 0100\rangle - 0111\rangle - 1000\rangle - 1011\rangle$
$ 1110\rangle$	$ 0101\rangle + 0110\rangle + 1001\rangle - 1010\rangle$
$ 1111\rangle$	$ 0101\rangle - 0110\rangle + 1001\rangle + 1010\rangle$

Since our scheme involves only CNOT [25], Hadamard [26] and controlled phase shift gates [27], it is completely feasible.

B. NDD of five qubit cluster states

There are 32 orthogonal $|C_5\rangle$ states which need to be distinguished for several quantum protocols. For this purpose, we make use of the five ancillas and make it interact with the five qubit cluster state as per Fig. 4. The outcome of the measurement on the ancillas and the corresponding cluster state are shown in the Table-II.

IV. ILLUSTRATION

A. Quantum dialogue using NDD of cluster states

We now devise a scheme for "quantum dialogue" using reusable cluster states with dense coding. The four and the five qubit cluster states reach the "Holevo bound" allowing four classical bits to be transmitted by sending two qubits and five classical bits be transmitted by sending three qubits respectively. In the standard dense coding protocol [4] involving four qubit cluster states, Alice and Bob possess the first and the last two qubits respectively. Alice encodes four classical bits of information into two qubits by performing a $(\sigma_i \otimes \sigma_j)$ operation where $i, j \in (0, 1, 2, 3)$. Alice, now sends her qubit to Bob. On receiving the qubits from Alice, Bob can perform our NDD scheme to decode the message without destroying $|C_4\rangle$. This is shown in Fig. 5.

TABLE II: Five qubit cluster state discrimination from the ancilla measurements

Ancilla Measurement	Corresponding five qubit Cluster State
00000⟩	00000⟩ + 00111⟩ + 11101⟩ + 11010⟩
00001⟩	00000⟩ - 00111⟩ + 11101⟩ - 11010⟩
00010⟩	00010⟩ + 00101⟩ + 11111⟩ - 11000⟩
00011⟩	00010⟩ - 00101⟩ + 11111⟩ - 11000⟩
00100⟩	00000⟩ + 00111⟩ - 11101⟩ - 11010⟩
00101⟩	00000⟩ - 00111⟩ - 11101⟩ + 11010⟩
00110⟩	00010⟩ + 00101⟩ - 11111⟩ - 11000⟩
00111⟩	00010⟩ - 00101⟩ - 11111⟩ + 11000⟩
01000⟩	01000⟩ + 01111⟩ + 10101⟩ + 10010⟩
01001⟩	01000⟩ - 01111⟩ + 10101⟩ - 10010⟩
01010⟩	01010⟩ + 01101⟩ + 10111⟩ + 10000⟩
01011⟩	01010⟩ - 01101⟩ + 10111⟩ - 10000⟩
01100⟩	01000⟩ + 01111⟩ - 10101⟩ - 10010⟩
01101⟩	01000⟩ - 01111⟩ - 10101⟩ + 10010⟩
01110⟩	01101⟩ + 01010⟩ - 10000⟩ - 10111⟩
01111⟩	01010⟩ - 01101⟩ - 10111⟩ + 10000⟩
10000⟩	00001⟩ + 00110⟩ + 11100⟩ + 11011⟩
10001⟩	00001⟩ - 00110⟩ + 11100⟩ - 11011⟩
10010⟩	00011⟩ + 00100⟩ + 11110⟩ + 11001⟩
10011⟩	00011⟩ - 00100⟩ + 11110⟩ - 11001⟩
10100⟩	00001⟩ + 00110⟩ - 11100⟩ - 11011⟩
10101⟩	00001⟩ - 00110⟩ - 11100⟩ + 11011⟩
10110⟩	00011⟩ + 00100⟩ - 11110⟩ - 11001⟩
10111⟩	00011⟩ - 00100⟩ - 11110⟩ + 11001⟩
11000⟩	01001⟩ + 01110⟩ + 10100⟩ + 10011⟩
11001⟩	01001⟩ - 01110⟩ + 10100⟩ - 10011⟩
11010⟩	01011⟩ + 01100⟩ + 10110⟩ + 10001⟩
11011⟩	01011⟩ - 01100⟩ + 10110⟩ - 10001⟩
11100⟩	01001⟩ + 01110⟩ - 10100⟩ - 10011⟩
11101⟩	01001⟩ - 01110⟩ - 10100⟩ + 10011⟩
11110⟩	01011⟩ + 01100⟩ - 10110⟩ - 10001⟩
11111⟩	01011⟩ - 01100⟩ - 10110⟩ + 10001⟩

Acknowledgments

The work was supported by the summer project programme undertaken by the Indian Institute of Science Education and Research (Kolkata), India. The authors acknowledge Sidharth K for discussions.

-
- [1] C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
 - [2] M. Hillery, V. Bužek, and A. Berthiaume, Phys. Rev. A **59**, 1829 (1999).
 - [3] A.M. Lance, T. Symul, W.P. Bowen, B.C. Sanders, and P.K. Lam, Phys. Rev. Lett. **92**, 177903 (2004).
 - [4] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. **69**, 2881 (1992).
 - [5] R. Raussendorf and H. J. Briegel, Phys. Rev. Lett. **86**, 5188 (2001).
 - [6] H. J. Briegel and R. Raussendorf, Phys. Rev. Lett. **86**, 910 (2001).
 - [7] C. Y. Lu, X. Q. Zhou, O. Ghne, W. B. Gao, J. Zhang, Z. S. Yuan, A. Goebel, T. Yang, and J. W. Pan, Nature **3**, 91 (2007).
 - [8] P. J. Blythe and B. T. H Varcoe, New. Jour. Phys. **8**, 231 (2006).
 - [9] Y. Soudagar, F. Bussières, G. Berln, S. Lacroix, J. M. Fernandez, and N. Godbout, J. Opt. Soc. Amer. B **24**, 226 (2007).
 - [10] M. Hein, W. Dur, and H. J. Briegel, Phys. Rev. A **71**, 032350 (2005).
 - [11] P. Walther, K. J. Resch, T. Rudolph, E. Schenck, H. Weinfurter, V. Vedral, M. Aspelmeyer, and A. Zeilinger, Nature **86**, 434 (169) 2005.
 - [12] Y. K. Bai and Z. D. Wang, eprint quant-ph/0709.4642v1.
 - [13] D. Liu, X. Zhao, and G. L. Long, eprint quant-ph/0705.3904v4.
 - [14] D. Schlingemann and R. F. Werner, Phys. Rev. A **65**, 012308 (2001).
 - [15] S. Muralidharan and P. K. Panigrahi, Phys. Rev. A **78**, 062333 (2008).
 - [16] S. Muralidharan and P. K. Panigrahi, Phys. Rev. A **77**, 032321 (2008).

- [17] S. Choudhury, S. Muralidharan and P. K. Panigrahi, J. Phys. A: Math. Gen. **42**, 115303 (2009).
- [18] Y. Yeo and W. K. Chua, Phys. Rev. Lett. **96**, 060502 (2006).
- [19] S. Muralidharan, S. Karumanchi, and P. K. Panigrahi, eprint quant-ph/0804.4206v2.
- [20] M. A. Nielsen, E. Knill, and R. Laflamme, Nature **96**, 52 (1998).
- [21] F. de Pasquale, G. Giorgi, and S. Paganelli, AIP. Conf. Proc. **734**, 48 (2004).
- [22] M. Gupta and P. K. Panigrahi, eprint quant-ph/0504183.
- [23] M. Gupta, A. Pathak, R. Srikanth, and P. K. Panigrahi, Int. Jour. Quant. Info. **5**, 627 (2007).
- [24] S. Muralidharan and P. K. Panigrahi, eprint quant-ph/0904.0563.
- [25] C. Monroe, D.M. Meekhof, B.E. King, W.M. Itano, and D.J. Wineland, Phys. Rev. Lett. **75**, 4714 (1995) .
- [26] V.I. Kuvshinov and A.V. Kuzmin, Phys. Lett. A **341**, 450 (2005).
- [27] T. Gopinath and A. Kumar, Jour. Mag. Res. **193**, 168 (2008).